



# The Eulerian Distribution on Involutions is Indeed Unimodal

Victor J. W. Guo, Jiang Zeng

## ► To cite this version:

Victor J. W. Guo, Jiang Zeng. The Eulerian Distribution on Involutions is Indeed Unimodal. 2006.  
hal-00018802

**HAL Id: hal-00018802**

**<https://hal.science/hal-00018802>**

Preprint submitted on 9 Feb 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# The Eulerian Distribution on Involutions is Indeed Unimodal

Victor J. W. Guo<sup>1</sup> and Jiang Zeng<sup>2</sup>

Institut Camille Jordan, Université Claude Bernard (Lyon I), F-69622, Villeurbanne Cedex, France

**Abstract.** Let  $I_{n,k}$  (respectively,  $J_{n,k}$ ) be the number of involutions (respectively, fixed-point free involutions) of  $\{1, \dots, n\}$  with  $k$  descents. Motivated by Brenti's conjecture which states that the sequence  $I_{n,0}, I_{n,1}, \dots, I_{n,n-1}$  is log-concave, we prove that the two sequences  $I_{n,k}$  and  $J_{2n,k}$  are unimodal in  $k$ , for all  $n$ . Furthermore, we conjecture that there are nonnegative integers  $a_{n,k}$  such that

$$\sum_{k=0}^{n-1} I_{n,k} t^k = \sum_{k=0}^{\lfloor (n-1)/2 \rfloor} a_{n,k} t^k (1+t)^{n-2k-1}.$$

This statement is stronger than the unimodality of  $I_{n,k}$  but is also interesting in its own right.

*Keywords:* Involutions, Descent number, Unimodality, Eulerian polynomial, Zeilberger's algorithm

*AMS Subject Classifications (2000):* Primary 05A15; Secondary 05A20

## 1 Introduction

A sequence  $a_0, a_1, \dots, a_n$  of real numbers is said to be *unimodal* if for some  $0 \leq j \leq n$  we have  $a_0 \leq a_1 \leq \dots \leq a_j \geq a_{j+1} \geq \dots \geq a_n$ , and is said to be *log-concave* if  $a_i^2 \geq a_{i-1}a_{i+1}$  for all  $1 \leq i \leq n-1$ . Clearly a log-concave sequence of *positive* terms is unimodal. The reader is referred to Stanley's survey [10] for the surprisingly rich variety of methods to show that a sequence is log-concave or unimodal. As noticed by Brenti [2], even though log-concave and unimodality have one-line definitions, to prove the unimodality or log-concavity of a sequence can sometimes be a very difficult task requiring the use of intricate combinatorial constructions or of refined mathematical tools.

Let  $\mathfrak{S}_n$  be the set of all permutations of  $[n] := \{1, \dots, n\}$ . We say that a permutation  $\pi = a_1 a_2 \dots a_n \in \mathfrak{S}_n$  has a *descent* at  $i$  ( $1 \leq i \leq n-1$ ) if  $a_i > a_{i+1}$ . The number of descents of  $\pi$  is called its descent number and is denoted by  $d(\pi)$ . A statistic on  $\mathfrak{S}_n$  is said to be *Eulerian*, if it is equidistributed with the descent number statistic. Recall that the polynomial

$$A_n(t) = \sum_{\pi \in \mathfrak{S}_n} t^{1+d(\pi)} = \sum_{k=1}^n A(n, k) t^k$$

is called an *Eulerian polynomial*. It is well-known that the *Eulerian numbers*  $A(n, k)$  ( $1 \leq k \leq n$ ) form a unimodal sequence, of which several proofs have been published: such

<sup>1</sup>guo@math.univ-lyon1.fr, <http://math.univ-lyon1.fr/~guo>

<sup>2</sup>zeng@math.univ-lyon1.fr, <http://math.univ-lyon1.fr/~zeng>

as the analytical one by showing that the polynomial  $A_n(t)$  has only real zeros [3, p. 294], by induction based on the recurrence relation of  $A(n, k)$  (see [9]), or by combinatorial techniques (see [7, 11]).

Let  $\mathcal{I}_n$  be the set of all involutions in  $\mathfrak{S}_n$  and  $\mathcal{J}_n$  the set of all fixed-point free involutions in  $\mathfrak{S}_n$ . Define

$$I_n(t) = \sum_{\pi \in \mathcal{I}_n} t^{d(\pi)} = \sum_{k=0}^{n-1} I_{n,k} t^k,$$

$$J_n(t) = \sum_{\pi \in \mathcal{J}_n} t^{d(\pi)} = \sum_{k=0}^{n-1} J_{n,k} t^k.$$

The first values of these polynomials are given in Table 1.

Table 1: The polynomials  $I_n(t)$  and  $J_n(t)$  for  $n \leq 6$ .

$n$	$I_n(t)$	$J_n(t)$
1	1	0
2	$1 + t$	$t$
3	$1 + 2t + t^2$	0
4	$1 + 4t + 4t^2 + t^3$	$t + t^2 + t^3$
5	$1 + 6t + 12t^2 + 6t^3 + t^4$	0
6	$1 + 9t + 28t^2 + 28t^3 + 9t^4 + t^5$	$t + 3t^2 + 7t^3 + 3t^4 + t^5$

As one may notice from Table 1 that the coefficients of  $I_n(t)$  and  $J_n(t)$  are *symmetric* and *unimodal* for  $1 \leq n \leq 6$ . Actually, the symmetries had been conjectured by Dumont and were first proved by Strehl [12]. Recently, Brenti (see [5]) conjectured that the coefficients of the polynomial  $I_n(t)$  are *log-concave* and Dukes [5] has obtained some partial results on the unimodality of the coefficients of  $I_n(t)$  and  $J_{2n}(t)$ . Note that, in contrast to Eulerian polynomials  $A_n(t)$ , the polynomials  $I_n(t)$  and  $J_{2n}(t)$  may have *non-real zeros*.

In this paper we will prove that for  $n \geq 1$ , the two sequences  $I_{n,0}, I_{n,1}, \dots, I_{n,n-1}$  and  $J_{2n,1}, J_{2n,2}, \dots, J_{2n,2n-1}$  are unimodal. Our starting point is the known generating functions of polynomials  $I_n(t)$  and  $J_n(t)$ :

$$\sum_{n=0}^{\infty} I_n(t) \frac{u^n}{(1-t)^{n+1}} = \sum_{r=0}^{\infty} \frac{t^r}{(1-u)^{r+1} (1-u^2)^{r(r+1)/2}}, \quad (1.1)$$

$$\sum_{n=0}^{\infty} J_n(t) \frac{u^n}{(1-t)^{n+1}} = \sum_{r=0}^{\infty} \frac{t^r}{(1-u^2)^{r(r+1)/2}}, \quad (1.2)$$

which have been obtained by Désarménien and Foata [4] and Gessel and Reutenauer [8] using different methods. We first derive linear recurrence formulas for  $I_{n,k}$  and  $J_{2n,k}$  in the next section and then prove the unimodality by induction in Section 3. We end this paper with further conjectures beyond the unimodality of the two sequences  $I_{n,k}$  and  $J_{2n,k}$ .

## 2 Linear recurrence formulas for $I_{n,k}$ and $J_{2n,k}$

Since the recurrence formula for the numbers  $I_{n,k}$  is a little more complicated than  $J_{2n,k}$ , we shall first prove it for the latter.

**Theorem 2.1.** *For  $n \geq 2$  and  $k \geq 0$ , the numbers  $J_{2n,k}$  satisfy the following recurrence formula:*

$$2nJ_{2n,k} = [k(k+1) + 2n - 2]J_{2n-2,k} + 2[(k-1)(2n-k-1) + 1]J_{2n-2,k-1} \\ + [(2n-k)(2n-k+1) + 2n - 2]J_{2n-2,k-2}. \quad (2.1)$$

Here and in what follows  $J_{2n,k} = 0$  if  $k < 0$ .

*Proof.* Equating the coefficients of  $u^{2n}$  in (1.2), we obtain

$$\frac{J_{2n}(t)}{(1-t)^{2n+1}} = \sum_{r=0}^{\infty} \binom{r(r+1)/2 + n - 1}{n} t^r. \quad (2.2)$$

Since

$$\binom{r(r+1)/2 + n - 1}{n} = \frac{r(r-1)/2 + r + n - 1}{n} \binom{r(r+1)/2 + n - 2}{n-1},$$

it follows from (2.2) that

$$\frac{J_{2n}(t)}{(1-t)^{2n+1}} = \frac{t^2}{2n} \left( \frac{J_{2n-2}(t)}{(1-t)^{2n-1}} \right)'' + \frac{t}{n} \left( \frac{J_{2n-2}(t)}{(1-t)^{2n-1}} \right)' + \frac{n-1}{n} \frac{J_{2n-2}(t)}{(1-t)^{2n-1}},$$

or

$$J_{2n}(t) = \frac{t^2(1-t)^2}{2n} J_{2n-2}''(t) + \left[ \frac{(2n-1)t^2(1-t)}{n} + \frac{t(1-t)^2}{n} \right] J_{2n-2}'(t) \\ + \left[ (2n-1)t^2 + \frac{(2n-1)(1-t)t}{n} + \frac{(n-1)(1-t)^2}{n} \right] J_{2n-2}(t) \\ = \frac{t^4 - 2t^3 + t^2}{2n} J_{2n-2}''(t) + \left[ \frac{(2-2n)t^3}{n} + \frac{(2n-3)t^2}{n} + \frac{t}{n} \right] J_{2n-2}'(t) \\ + \left[ (2n-2)t^2 + \frac{t}{n} + \frac{n-1}{n} \right] J_{2n-2}(t). \quad (2.3)$$

Equating the coefficients of  $t^n$  in (2.3) yields

$$J_{2n,k} = \frac{(k-2)(k-3)}{2n} J_{2n-2,k-2} - \frac{(k-1)(k-2)}{n} J_{2n-2,k-1} + \frac{k(k-1)}{2n} J_{2n-2,k} \\ + \frac{(2-2n)(k-2)}{n} J_{2n-2,k-2} + \frac{(2n-3)(k-1)}{n} J_{2n-2,k-1} + \frac{k}{n} J_{2n-2,k} \\ + (2n-2)J_{2n-2,k-2} + \frac{1}{n} J_{2n-2,k-1} + \frac{n-1}{n} J_{2n-2,k}.$$

After simplification, we obtain (2.1). ■

**Theorem 2.2.** For  $n \geq 3$  and  $k \geq 0$ , the numbers  $I_{n,k}$  satisfy the following recurrence formula:

$$\begin{aligned} nI_{n,k} &= (k+1)I_{n-1,k} + (n-k)I_{n-1,k-1} + [(k+1)^2 + n-2]I_{n-2,k} \\ &\quad + [2k(n-k-1) - n+3]I_{n-2,k-1} + [(n-k)^2 + n-2]I_{n-2,k-2}. \end{aligned} \quad (2.4)$$

Here and in what follows  $I_{n,k} = 0$  if  $k < 0$ .

*Proof.* Extracting the coefficients of  $u^{2n}$  in (1.1), we obtain

$$\frac{I_n(t)}{(1-t)^{n+1}} = \sum_{r=0}^{\infty} t^r \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{r(r+1)/2 + k - 1}{k} \binom{r+n-2k}{n-2k}. \quad (2.5)$$

Let

$$T(n, k) := \binom{x+k-1}{k} \binom{y+n-2k}{n-2k},$$

and

$$s(n) := \sum_{k=0}^{\lfloor n/2 \rfloor} T(n, k).$$

Applying Zeilberger's algorithm, the Maple package `ZeilbergerRecurrence(T,n,k,s,0..n)` gives

$$(2x+y+n+1)s(n) + (y+1)s(n+1) - (n+2)s(n+2) = 0, \quad (2.6)$$

i.e.,

$$s(n) = \frac{y+1}{n}s(n-1) + \frac{2x+y+n-1}{n}s(n-2).$$

When  $x = r(r+1)/2$  and  $y = r$ , we get

$$s(n) = \frac{r+1}{n}s(n-1) + \frac{r(r-1) + 3r + n - 1}{n}s(n-2). \quad (2.7)$$

Now, from (2.5) and (2.7) follows the identity

$$\begin{aligned} \frac{nI_n(t)}{(1-t)^{n+1}} &= t \left( \frac{I_{n-1}(t)}{(1-t)^n} \right)' + \frac{I_{n-1}(t)}{(1-t)^n} + t^2 \left( \frac{I_{n-2}(t)}{(1-t)^{n-1}} \right)'' + 3t \left( \frac{I_{n-2}(t)}{(1-t)^{n-1}} \right)' \\ &\quad + (n-1) \frac{I_{n-2}(t)}{(1-t)^{n-1}}, \end{aligned}$$

or

$$\begin{aligned} nI_n(t) &= (t-t^2)I'_{n-1}(t) + [1+(n-1)t]I_{n-1}(t) + t^2(1-t)^2I''_{n-2}(t) \\ &\quad + t(1-t)[3+(2n-5)t]I'_{n-2}(t) + (n-1)[1+t+(n-2)t^2]I_{n-2}(t). \end{aligned} \quad (2.8)$$

Comparing the coefficients of  $t^k$  in both sides of (2.8), we obtain

$$\begin{aligned} nI_{n,k} &= kI_{n-1,k} - (k-1)I_{n-1,k-1} + I_{n-1,k} + (n-1)I_{n-1,k-1} \\ &\quad + k(k-1)I_{n-2,k} - 2(k-1)(k-2)I_{n-2,k-1} + (k-2)(k-3)I_{n-2,k-2} \\ &\quad + 3kI_{n-2,k} + (2n-8)(k-1)I_{n-2,k-1} - (2n-5)(k-2)I_{n-2,k-2} \\ &\quad + (n-1)I_{n-2,k} + (n-1)I_{n-2,k-1} + (n-1)(n-2)I_{n-2,k-2}, \end{aligned}$$

which, after simplification, equals the right-hand side of (2.4). ■

*Remark.* The recurrence formula (2.6) can also be proved by hand as follows. It is easy to see that the generating function of  $s(n)$  is

$$\sum_{n=0}^{\infty} s(n)u^n = (1-u^2)^{-x}(1-u)^{-y-1}. \quad (2.9)$$

Differentiating (2.9) with respect to  $u$  implies that

$$\sum_{n=0}^{\infty} ns(n)u^{n-1} = \left( \frac{2ux}{1-u^2} + \frac{y+1}{1-u} \right) (1-u^2)^{-x}(1-u)^{-y-1},$$

consequently,

$$\begin{aligned} (1-u^2) \sum_{n=0}^{\infty} ns(n)u^{n-1} &= [(2x+y+1)u + y+1](1-u^2)^{-x}(1-u)^{-y-1} \\ &= [(2x+y+1)u + y+1] \sum_{n=0}^{\infty} s(n)u^n. \end{aligned} \quad (2.10)$$

Comparing the coefficients of  $u^{n+1}$  in both sides of (2.10), we obtain

$$(n+2)s(n+2) - ns(n) = (2x+y+1)s(n) + (y+1)s(n+1),$$

which is equivalent to (2.6).

Note that the right-hand side of (2.1) (respectively, (2.4)) is invariant under the substitution  $k \rightarrow 2n-k$  (respectively,  $k \rightarrow n-1-k$ ), provided that the sequence  $I_{n-1,k}$  (respectively,  $J_{2n-2,k}$ ) is symmetric. Thus, by induction we derive immediately the symmetry properties of  $J_{2n,k}$  and  $I_{n,k}$  (see [4, 8, 12]).

**Corollary 2.3.** *For  $n, k \in \mathbb{N}$ , we have*

$$I_{n,k} = I_{n,n-1-k}, \quad J_{2n,k} = J_{2n,2n-k}.$$

It would be interesting to find a combinatorial proof of the recurrence formulas (2.1) and (2.4), since such a proof could hopefully lead to a combinatorial proof of the unimodality of these two sequences.

### 3 Unimodality of the sequences $I_{n,k}$ and $J_{2n,k}$

The following observation is crucial in our inductive proof of the unimodality of the sequences  $I_{n,k}$  ( $0 \leq k \leq n-1$ ) and  $J_{2n,k}$  ( $1 \leq k \leq 2n-1$ ).

**Lemma 3.1.** *Let  $x_0, x_1, \dots, x_n$  and  $a_0, a_1, \dots, a_n$  be real numbers such that  $x_0 \geq x_1 \geq \dots \geq x_n \geq 0$  and  $a_0 + a_1 + \dots + a_k \geq 0$  for all  $k = 0, 1, \dots, n$ . Then*

$$\sum_{i=0}^n a_i x_i \geq 0.$$

Indeed, the above inequality follows from the identity:

$$\sum_{i=0}^n a_i x_i = \sum_{k=0}^n (x_k - x_{k+1})(a_0 + a_1 + \dots + a_k),$$

where  $x_{n+1} = 0$ .

**Theorem 3.2.** *The sequence  $J_{2n,1}, J_{2n,2}, \dots, J_{2n,2n-1}$  is unimodal.*

*Proof.* By the symmetry of  $J_{2n,k}$ , it is enough to show that  $J_{2n,k} \geq J_{2n,k-1}$  for all  $2 \leq k \leq n$ . We proceed by induction on  $n$ . The  $n = 2$  case is clear from Table 1. Suppose the sequence  $J_{2n-2,k}$  is unimodal in  $k$ . By Theorem 2.1, one has

$$2n(J_{2n,k} - J_{2n,k-1}) = A_0 J_{2n-2,k} + A_1 J_{2n-2,k-1} + A_2 J_{2n-2,k-2} + A_3 J_{2n-2,k-3}, \quad (3.1)$$

where

$$\begin{aligned} A_0 &= k^2 + k + 2n - 2, & A_1 &= 4nk - 3k^2 - 6n + k + 6, \\ A_2 &= 3k^2 + 4n^2 - 8nk - 5k + 12n - 4, & A_3 &= 3k - k^2 + 4nk - 4n^2 - 8n. \end{aligned}$$

We have the following two cases:

- If  $2 \leq k \leq n-1$ , then

$$J_{2n-2,k} \geq J_{2n-2,k-1} \geq J_{2n-2,k-2} \geq J_{2n-2,k-3}$$

by the induction hypothesis, and clearly

$$\begin{aligned} A_0 &\geq 0, & A_0 + A_1 &= 2(k-1)(2n-k) + 4 \geq 0, \\ A_0 + A_1 + A_2 &= (2n-k)^2 - 3k + 8n \geq 0, & A_0 + A_1 + A_2 + A_3 &= 0. \end{aligned}$$

Therefore, by Lemma 3.1, we have

$$J_{2n,k} - J_{2n,k-1} \geq 0.$$

- If  $k = n$ , then

$$J_{2n-2,n-1} \geq J_{2n-2,n} = J_{2n-2,n-2} \geq J_{2n-2,n-3}$$

by symmetry and the induction hypothesis. In this case, we have  $A_1 = (n-2)(n-3) \geq 0$  and thus the corresponding condition of Lemma 3.1 is satisfied. Therefore, we have

$$J_{2n,n} - J_{2n,n-1} \geq 0.$$

This completes the proof. ■

**Theorem 3.3.** *The sequence  $I_{n,0}, I_{n,1}, \dots, I_{n,n-1}$  is unimodal.*

*Proof.* By the symmetry of  $I_{n,k}$ , it suffices to show that  $I_{n,k} \geq I_{n,k-1}$  for all  $1 \leq k \leq (n-1)/2$ . From Table 1, it is clear that the sequences  $I_{n,k}$  are unimodal in  $k$  for  $1 \leq n \leq 6$ .

Now suppose  $n \geq 7$  and the sequences  $I_{n-1,k}$  and  $I_{n-2,k}$  are unimodal in  $k$ . Replacing  $k$  by  $k-1$  in (2.4), we obtain

$$\begin{aligned} nI_{n,k-1} &= kI_{n-1,k-1} + (n-k+1)I_{n-1,k-2} + (k^2 + n-2)I_{n-2,k-1} \\ &\quad + [2(k-1)(n-k) - n+3]I_{n-2,k-2} + [(n-k+1)^2 + n-2]I_{n-2,k-3}. \end{aligned} \quad (3.2)$$

Combining (2.4) and (3.2) yields

$$\begin{aligned} n(I_{n,k} - I_{n,k-1}) &= B_0I_{n-1,k} + B_1I_{n-1,k-1} + B_2I_{n-1,k-2} \\ &\quad + C_0I_{n-2,k} + C_1I_{n-2,k-1} + C_2I_{n-2,k-2} + C_3I_{n-2,k-3}, \end{aligned} \quad (3.3)$$

where

$$\begin{aligned} B_0 &= k+1, \quad B_1 = n-2k, \quad B_2 = -(n-k+1), \\ C_0 &= (k+1)^2 + n-2, \quad C_1 = 2nk - 3k^2 - 2k - 2n + 5, \\ C_2 &= n^2 - 4nk + 3k^2 + 4n - 2k - 5, \quad C_3 = -(n-k+1)^2 - n + 2. \end{aligned}$$

Notice that  $I_{n-1,k} \geq I_{n-1,k-1} \geq I_{n-1,k-2}$  for  $1 \leq k \leq (n-1)/2$ . By Lemma 3.1, we have

$$B_0I_{n-1,k} + B_1I_{n-1,k-1} + B_2I_{n-1,k-2} \geq 0. \quad (3.4)$$

It remains to show that

$$C_0I_{n-2,k} + C_1I_{n-2,k-1} + C_2I_{n-2,k-2} + C_3I_{n-2,k-3} \geq 0, \quad \forall 1 \leq k \leq (n-1)/2. \quad (3.5)$$

We need to consider the following two cases:

- If  $1 \leq k \leq (n-2)/2$ , then

$$I_{n-2,k} \geq I_{n-2,k-1} \geq I_{n-2,k-2} \geq I_{n-2,k-3}$$

by the induction hypothesis, and

$$\begin{aligned} C_0 &= (k+1)^2 + n-2 \geq 0, \quad C_0 + C_1 = (2k-1)(n-k-1) + k+3 \geq 0, \\ C_0 + C_1 + C_2 &= (n-k+1)^2 + n-2 \geq 0, \quad C_0 + C_1 + C_2 + C_3 = 0. \end{aligned}$$



- If  $k = (n - 1)/2$ , then by symmetry and the induction hypothesis,

$$I_{n-2,k-1} \geq I_{n-2,k} = I_{n-2,k-2} \geq I_{n-2,k-3}.$$

In this case, we have  $C_1 = (n - 3)(n - 7)/4 \geq 0$  for  $n \geq 7$ .

Therefore, by Lemma 3.1 the inequality (3.5) holds. It follows from (3.3)–(3.5) that

$$I_{n,k} - I_{n,k-1} \geq 0, \quad \forall 1 \leq k \leq (n - 1)/2.$$

This completes the proof. ■

## 4 Further remarks and open problems

Since  $I_{n,k} = I_{n,n-1-k}$ , we can rewrite  $I_n(t)$  as follows:

$$I_n(t) = \sum_{k=0}^{n-1} I_{n,k} t^k = \begin{cases} \sum_{k=0}^{n/2-1} I_{n,k} t^k (1 + t^{n-2k-1}), & \text{if } n \text{ is even,} \\ I_{n,(n-1)/2} t^{(n-1)/2} + \sum_{k=0}^{(n-3)/2} I_{n,k} t^k (1 + t^{n-2k-1}), & \text{if } n \text{ is odd.} \end{cases}$$

Applying the well-known formula

$$x^n + y^n = \sum_{j=0}^{\lfloor n/2 \rfloor} (-1)^j \frac{n}{n-j} \binom{n-j}{j} (xy)^j (x+y)^{n-2j},$$

we obtain

$$I_n(t) = \sum_{k=0}^{\lfloor (n-1)/2 \rfloor} a_{n,k} t^k (1+t)^{n-2k-1}, \quad (4.1)$$

where

$$a_{n,k} = \begin{cases} \sum_{j=0}^k (-1)^{k-j} \frac{n-2j-1}{n-k-j-1} \binom{n-k-j-1}{k-j} I_{n,j}, & \text{if } 2k+1 < n, \\ I_{n,k} + \sum_{j=0}^{k-1} (-1)^{k-j} \frac{n-2j-1}{n-k-j-1} \binom{n-k-j-1}{k-j} I_{n,j}, & \text{if } 2k+1 = n. \end{cases}$$

The first values of  $a_{n,k}$  are given in Table 2, which seems to suggest the following conjecture.

**Conjecture 4.1.** *For  $n \geq 1$  and  $k \geq 0$ , the coefficients  $a_{n,k}$  are nonnegative integers.*

Table 2: Values of  $a_{n,k}$  for  $n \leq 16$  and  $0 \leq k \leq \lfloor (n-1)/2 \rfloor$ .

$k \setminus n$	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
1			0	1	2	4	6	9	12	16	20	25	30	36	42	49
2					2	6	18	39	79	141	239	379	579	849	1211	1680
3							0	18	78	272	722	1716	3626	7160	13206	23263
4									20	124	668	2560	8360	23536	59824	139457
5											32	700	4800	24160	95680	325572
6													440	5480	44632	257964
7															2176	44376

Since the coefficients of  $t^k(1+t)^{n-2k-1}$  are symmetric and unimodal with center of symmetry at  $(n-1)/2$ , Conjecture 4.1 is stronger than the fact that the coefficients of  $I_n(t)$  are symmetric and unimodal. A more interesting question is to give a combinatorial interpretation of  $a_{n,k}$ . Note that the Eulerian polynomials can be written as

$$A_n(t) = \sum_{k=1}^{\lfloor (n+1)/2 \rfloor} c_{n,k} t^k (1+t)^{n-2k+1},$$

where  $c_{n,k}$  is the number of increasing binary trees on  $[n]$  with  $k$  leaves and no vertices having left children only (see [1, 6, 7]).

We now proceed to derive a recurrence relation for  $a_{n,k}$ . Set  $x = x(t) = t/(1+t)^2$  and

$$P_n(x) = \sum_{k=0}^{\lfloor (n-1)/2 \rfloor} a_{n,k} x^k.$$

Then we can rewrite (4.1) as

$$I_n(t) = (1+t)^{n-1} P_n(x). \quad (4.2)$$

Differentiating (4.2) with respect to  $t$  we get

$$I'_n(t) = (n-1)(1+t)^{n-2} P_n(x) + (1+t)^{n-1} P'_n(x) x'(t), \quad (4.3)$$

$$\begin{aligned} I''_n(t) &= (n-1)(n-2)(1+t)^{n-3} P_n(x) + 2(n-1)(1+t)^{n-2} P'_n(x) x'(t) \\ &\quad + (1+t)^{n-1} P''_n(x) (x'(t))^2 + (1+t)^{n-1} P'_n(x) x''(t), \end{aligned} \quad (4.4)$$

$$x'(t) = \frac{1-t}{(1+t)^3}, \quad x''(t) = \frac{2t-4}{(1+t)^4}. \quad (4.5)$$

Substituting (4.2)–(4.5) into (2.8), we obtain

$$\begin{aligned}
& n(1+t)^{n-1}P_n(x) \\
&= [1 + (2n-2)t + t^2](1+t)^{n-3}P_{n-1}(x) + t(1-t)^2(1+t)^{n-5}P'_{n-1}(x) \\
&\quad + [-(t^2 + 14t + 1)(1-t)^2 + (1 + 6t - 18t^2 + 6t^3 + t^4)n + 4t^2n^2](1+t)^{n-5}P_{n-2}(x) \\
&\quad + [3t(t^2 - 4t + 1)(1-t)^2 + 4t^2(1-t)^2n](1+t)^{n-7}P'_{n-2}(x) \\
&\quad + t^2(1-t)^4(1+t)^{n-9}P''_{n-2}(x).
\end{aligned} \tag{4.6}$$

Dividing the two sides of (4.6) by  $(1+t)^{n-1}$  and noticing that  $t/(1+t)^2 = x$ , after a little manipulation we get

$$\begin{aligned}
nP_n(x) &= [1 + (2n-4)x]P_{n-1}(x) + (x-4x^2)P'_{n-1}(x) \\
&\quad + [(n-1) + (2n-8)x + 4(n-3)(n-4)x^2]P_{n-2}(x) \\
&\quad + [3x + (4n-30)x^2 + (72-16n)x^3]P'_{n-2}(x) + (x^2 - 8x^3 + 16x^4)P''_{n-2}(x).
\end{aligned}$$

Extracting the coefficients of  $x^k$  yields

$$\begin{aligned}
na_{n,k} &= a_{n-1,k} + (2n-4)a_{n-1,k-1} + ka_{n-1,k} - 4(k-1)a_{n-1,k-1} \\
&\quad + (n-1)a_{n-2,k} + (2n-8)a_{n-2,k-1} + 4(n-3)(n-4)a_{n-2,k-2} \\
&\quad + 3ka_{n-2,k} + (4n-30)(k-1)a_{n-2,k-1} + (72-16n)(k-2)a_{n-2,k-2} \\
&\quad + k(k-1)a_{n-2,k} - 8(k-1)(k-2)a_{n-2,k-1} + 16(k-2)(k-3)a_{n-2,k-2}.
\end{aligned}$$

After simplification, we obtain the following recurrence formula for  $a_{n,k}$ .

**Theorem 4.2.** *For  $n \geq 3$  and  $k \geq 0$ , there holds*

$$\begin{aligned}
na_{n,k} &= (k+1)a_{n-1,k} + (2n-4k)a_{n-1,k-1} + [k(k+2) + n-1]a_{n-2,k} \\
&\quad + [(k-1)(4n-8k-14) + 2n-8]a_{n-2,k-1} + 4(n-2k)(n-2k+1)a_{n-2,k-2},
\end{aligned} \tag{4.7}$$

where  $a_{n,k} = 0$  if  $k < 0$  or  $k > (n-1)/2$ .

Note that, if  $n \geq 2k+3$ , then

$$(k-1)(4n-8k-14) + 2n-8 > 0 \quad \text{for any } k \geq 1,$$

and so are the other coefficients in (4.7). Therefore, Conjecture 4.1 would be proved if one can show that  $a_{2n+1,n} \geq 0$  and  $a_{2n+2,n} \geq 0$ .

Finally, from (4.1) it is easy to see that

$$a_{2n+1,n} = (-1)^n I_{2n+1}(-1) = \sum_{k=0}^{2n} (-1)^{n-k} I_{2n+1,k},$$

$$a_{2n+2,n} = (-1)^n I'_{2n+2}(-1) = \sum_{k=1}^{2n+1} (-1)^{n+1-k} k I_{2n+2,k}.$$

Thus, Conjecture 4.1 is equivalent to the *nonnegativity* of the above two alternating sums.

Since  $J_{2n,k} = J_{2n,2n-k}$ , in the same manner as  $I_n(t)$  we obtain

$$J_{2n}(t) = \sum_{k=1}^n b_{2n,k} t^k (1+t)^{2n-2k},$$

where

$$b_{2n,k} = \begin{cases} \sum_{j=1}^k (-1)^{k-j} \frac{2n-2j}{2n-k-j} \binom{2n-k-j}{k-j} J_{2n,j}, & \text{if } k < n, \\ J_{2n,k} + \sum_{j=1}^{k-1} (-1)^{k-j} \frac{2n-2j}{2n-k-j} \binom{2n-k-j}{k-j} J_{2n,j}, & \text{if } k = n. \end{cases}$$

Now, it follows from (2.2) that

$$J_{2n,k} = \sum_{i=0}^k (-1)^{k-i} \binom{2n+1}{k-i} \binom{i(i+1)/2 + n - 1}{i(i+1)/2 - 1}$$

is a polynomial in  $n$  of degree  $d := k(k+1)/2 - 1$  with leading coefficient  $1/d!$ , and so is  $b_{2n,k}$ . Thus, we have  $\lim_{n \rightarrow +\infty} b_{2n,k} = +\infty$  for any fixed  $k > 1$ .

The first values of  $b_{2n,k}$  are given in Table 3, which seems to suggest

**Conjecture 4.3.** *For  $n \geq 9$  and  $k \geq 1$ , the coefficients  $b_{2n,k}$  are nonnegative integers.*

Similarly to the proof of Theorem 4.2, we can prove the following result.

**Theorem 4.4.** *For  $n \geq 2$  and  $k \geq 1$ , there holds*

$$2nb_{2n,k} = [k(k+1) + 2n - 2]b_{2n-2,k} + [2 + 2(k-1)(4n-4k-3)]b_{2n-2,k-1} \\ + 8(n-k+1)(2n-2k+1)b_{2n-2,k-2}.$$

where  $b_{2n,k} = 0$  if  $k < 1$  or  $k > n$ .

Theorem 4.4 allows us to reduce the verification of Conjecture 4.3 to the boundary case  $b_{2n,n} \geq 0$  for  $n \geq 9$ .

**Acknowledgment.** The second author was supported by EC's IHRP Programme, within Research Training Network "Algebraic Combinatorics in Europe," grant HPRN-CT-2001-00272.

Table 3: Values of  $b_{2n,k}$  for  $2n \leq 24$  and  $1 \leq k \leq n$ .

$k \setminus 2n$	2	4	6	8	10	12	14	16	18	20	22	24
1	1	1	1	1	1	1	1	1	1	1	1	1
2		-1	-1	0	2	5	9	14	20	27	35	44
3			3	12	36	91	201	399	728	1242	2007	3102
4				-7	-10	91	652	2593	7902	20401	46852	98494
5					25	219	1710	10532	50165	194139	639968	1861215
6						-65	249	11319	122571	841038	4377636	18747924
7							283	6586	135545	1737505	15219292	101116704
8								-583	33188	1372734	24412940	277963127
9									4417	379029	16488999	367507439
10										1791	3350211	203698690
11											133107	36903128
12												761785

## References

- [1] P. Brändén, Sign-graded posets, unimodality of  $W$ -polynomials and the Charney-Davis conjecture, *Electron. J. Combin.* **11** (2) (2004/05), #R9.
- [2] F. Brenti, Log-concave and unimodal sequences in algebra, combinatorics, and geometry: an update, *Jerusalem combinatorics'93*, pp. 71–89, *Contemp. Math.*, **178**, Amer. Math. Soc., Providence, RI, 1994.
- [3] L. Comtet, *Advanced Combinatorics*, Reidel, Dordrecht/Boston, 1974.
- [4] J. Désarménien and D. Foata, Fonctions symétriques et séries hypergéométriques basiques multivariées, *Bull. Soc. Math. France* **113** (1985), 3–22.
- [5] W. M. B. Dukes, Permutation statistics on involutions, *European J. Combin.*, to appear.
- [6] D. Foata and V. Strehl, Euler numbers and variations of permutations, *Atti dei Convegni Lincei*, **17** Tomo I, 1976, pp. 119–131.
- [7] V. Gasharov, On the Neggers-Stanley conjecture and the Eulerian polynomials, *J. Combin. Theory Ser. A* **82** (1998), 134–146.
- [8] I. M. Gessel and C. Reutenauer, Counting permutations with given cycle structure and descent set, *J. Combin. Theory Ser. A* **64** (1993), 189–215.
- [9] D. C. Kurtz, A note on concavity properties of triangular arrays of numbers, *J. Combin. Theory Ser. A* **13** (1972), 135–139.
- [10] R. P. Stanley, Log-concave and unimodal sequences in algebra, combinatorics, and geometry, *Ann. New York Acad. Sci.* **576** (1989), 500–535.
- [11] J. R. Stembridge, Eulerian numbers, tableaux, and the Betti numbers of a toric variety, *Discrete Math.* **99** (1992), 307–320.
- [12] V. Strehl, Symmetric Eulerian distributions for involutions, *Séminair Lotharingien Combinatoire* **1**, Strasbourg 1980, Publications de l'I.R.M.A. 140/S-02, Strasbourg, 1981.